

STRATIGRAPHY AND TECTONICS OF THE UPPERMOST BUCEGI CONGLOMERATE FORMATION (ALBIAN, EASTERN CARPATHIANS, ROMANIA)

DAN CONSTATIN JIPA, CONSTANTIN UNGUREANU AND GABRIEL ION

*National Institute of Marine Geology and Geo-ecology, 23-25 Dimitrie Onciul Street, RO-024053 Bucharest, Romania,
e-mail: jipa@geoecomar.ro, costinungureanu@geoecomar.ro, gion@geoecomar.ro*

Abstract. This paper presents a detailed lithostratigraphic investigation carried out in the Bucegi Massif, a mountainous zone from the southern part of the Eastern Carpathians. The study focused on the 50 - 70 m thick succession of four conglomerate units associated with olistoliths, intercalated in the upper part of the Babele Sandstone Member (the uppermost lithostratigraphic unit of the Albian Bucegi Conglomerate Formation). The conglomerate beds succession stands for the uppermost strata of the thick Bucegi conglomeratic accumulation. All the four conglomerate bodies are stratiform and continuous for almost two kilometers, and show mainly constant lithofacies features on wide areas. The study emphasizes two large-scale trends of high implication on depositional interpretation. The reduced areal extension of the basal conglomerate unit, compared with the larger extension area of the overlying units 2, 3 and 4, is one of the features with genetic significance. Another remarkable characteristic is the lateral passage from the single thick conglomerate units 2, 3 and 4, to conglomerate units alternating with sandstones beds. Another remarkable characteristic is the northwards amalgamation of the conglomerate units 1, 2, 3 and 4 into a single body, named in this paper the Conglomerates with Olistoliths.

Key words: Bucegi Massif, Albian, lithostratigraphy, conglomerate unit, olistoliths, fining upward trend.

1. INTRODUCTION

The Bucegi Conglomerate Formation is the lithostratigraphic term referring to thick conglomerate deposits accumulated at the end of the Lower Cretaceous, cropping out in the Bucegi Massif area, at the southern extremity of the Eastern Carpathians.

Recent observations pointed out to a succession of several distinct Albian conglomerate units genetically related, but with unclear stratigraphic significance. In order to proceed to upcoming sedimentology interpretations, it was essential to build up an up-to-date model of the geologic architecture in the northern Bucegi study area. This became the goal of a detailed stratigraphy and tectonics study that is herein presented. Our investigation was focused at the level of the upper member of the Albian Bucegi Formation, namely the Babele Sandstone Member.

The present study was carried out in the Obârșia Mountain zone, between Șugărilor and Obârșiei Rivers (Fig. 1). This area is located in the northern part of the Bucegi Massif, where the map published by Patrușiu (1969) shows multiple olistoliths (Fig. 2). Geographically, the study area is situated in the Prahova County, in the spring area of the Ialomița River.

The Obârșia Mountain has two topography summits, which will be frequently referred to in this paper: the Colții Obârșiei Peak (2488 m) to the north, and a nameless summit with 2380 m elevation to the south (Fig. 1).

Geologic mapping was the main approach to accomplish the study objective. The map previously made by Patrușiu (1969) served as a starting point, the new cartographic activity looking for more detailed geologic features. Besides the classical rock-outcrop examination, special consideration was paid to the relief morphology, capable to reveal the trend of some geological boundaries, mainly in the areas with poor quality outcrop coverage. The satellite image examination,




Fig. 1. Areal location of the Obârșia Mountain geology investigations. The study area location is shown on the Romania index map (left), Bucegi Massif simplified geological map (based on Patrușiu, 1969) (middle) and on the Obârșia Mt. topographic map (right).

using Bing Map imagery, provided more arguments to the geologic boundaries mapping, and offered a detailed view to some critical features.

2. GEOLOGIC SETTING


2.1. OVERVIEW OF THE LITHOSTRATIGRAPHY IN THE BUCEGI MASSIF

The present stratigraphy outline of the Bucegi conglomeratic deposits was achieved only subsequent to the geologic study of the underlying Lower Cretaceous and overlying Upper Cretaceous deposits. The investigation of the marly-sandy flysch deposits from the Eastern Carpathians bend zone, which includes the Bucegi area, revealed the existence of relevant Aptian-Lower Albian macrofaunal assemblages. Murgeanu *et al.* (1963) present a list of fauna, including the following taxa: *Deshayesites borowae* (from Cărpiniș River), *Dufrenoya furcata*, *Colombiceras subpeltoceratoides*, *C. aff. tableri discoidalis*, *C. aff. crassicostatum*, and *Holcophylloceras* sp. (Valea Seacă River from Caraiman Mountain), *Leymeriella aff. revili* (Bușteni), *C. sp. tableri discoidalis* and *Parahoplites gr. grossouvrei* (Runcu River). Recent studies (Melinte and Jipa, 2007; Briceag *et al.*, 2009) identified at the top of the flysch deposits underlying the Bucegi Formation, the first occurrence of the nannofossil *Prediscosphaera columnata*, an Early Albian bioevent (Roth, 1983; Mutterlose, 1992), its age being 112.6 Ma (Bralower *et al.*, 1997).

As concerning the deposits covering the Bucegi Formation, Murgeanu and Patrușiu (1957), confirmed by Avram (1967), remarked that the Vraconian (*i.e.*, latest Albian) sediments are unconformably overlying conglomerates and sandstones of the Bucegi Formation from the Dâmbovița River area (Cotenești, Lăicăi and Bădeni localities). The Vraconian fauna of these deposits includes *Aucellina gryphaeoides*, *Parahibolites tourtiaie* and *Puzosia subplanulata*.

So far no fossils useful for the geologic age determination were discovered in the deposits of the Bucegi Formation. This thick conglomerate deposit was assigned to a Middle and Late Albian age by Murgeanu and Patrușiu (1963), taking into account their geometric position between the Aptian-Lower Albian flysch deposits and the Vraconian hemipelagic marls (Fig. 3). Subsequently, Patrușiu (1969) refers to the underlying flysch deposits as not younger than the Late Aptian, while the main Bucegi Conglomerate series was assigned to the Albian.

Within the conglomerate succession of the Bucegi Massif, Murgeanu and Patrușiu (1963) separated three main lithostratigraphic entities. At the base of this conglomerate accumulation are the *Lower Bucegi Conglomerates*, located in the upper part of the Aptian flysch deposits. The *Middle Bucegi Conglomerate* unit includes the major part of the Albian conglomerate assemblage. They are conformably covered by the *Upper Bucegi Conglomerate* unit, distinguished by the presence of embedded blocks and olistoliths.



1000 m




Fig 2. Study area location on the geological map of northern Bucegi Massif. From Patrulea *et al.* (1971) (geological map 1:50,000 scale, sheet 110D Moeciu). The geological map is draped on the 3D relief. Due to the perspective view, the graphic scale is approximate, mostly valid on the E-W direction.

In the lithostratigraphic column published by Murgeanu and Patrulea (1963) (Fig. 3), the upper part of the Albian Bucegi Formation is relatively finer-grained, with sandstones and microconglomerates. Patrulea (1969) described these deposits as the *Babele Sandstone*, which is the upper member of the Bucegi Formation (Fig. 4). The *Scropoasa-Lăptici Sandstone* is another sandy lithofacies unit from the Bucegi Massif, described by Patrulea (1969). Made of fine/silty sandstones with intercalated coarser-grained sandstone beds, the Scropoasa-Lăptici Sandstone unit occurs between the Middle and Upper Bucegi Conglomerates and thins out and disappears eastward (Fig. 4).

2.2. TECTONIC FEATURES OF THE ALBIAN DEPOSITS FROM THE NORTHERN BUCEGI MASSIF AREA

The Bucegi Formation represents the youngest Lower Cretaceous deposits of the Ceahlău Nappe unit. This is the main tectonic unit of the Outer Dacid Nappe System (Murgeanu *et al.*, 1963; Dumitrescu and Săndulescu, 1968; Săndulescu, 1984), with a large spatial development.

The tectonic setting of the Bucegi Conglomerate Formation was outlined by Murgeanu and Patrulea (1963) and corroborated by Murgeanu *et al.* (1963) and Patrulea (1969). In their view, the Albian sedimentation in the Eastern Carpathian bend area (including the Bucegi Massif) developed during the Mesocretaceous tectonic movements. As Patrulea

(1969) emphasized, since its early Aptian stage (end of the Bedoulian) up to its late Albian stage (pre-Vraconian), the Austrian Mesocretaceous diastrophism produced uplift and subsidence in different areas, and the emerged territory suffered strong erosion.




Fig. 3. Murgeanu *et al.* (1963) lithostratigraphic column of the Bucegi Conglomerate Formation. This is the first lithostratigraphic column of the entire conglomeratic accumulation from the Bucegi Massif.




Fig. 4. Patrulea view on the Bucegi Conglomerate sedimentary architecture. **A.** Geological section in the northern part of the Bucegi Massif. Redrawn after Patrulea (1969). **B.** Geological section in the median part of the Bucegi Massif. From Patrulea *et al.* (1971). **Legend:** 1. Crystalline schists (smuclab = muscovite-chlorite schists with porphyroblastic albite). 2. Middle Jurassic (Bajocian-Callovian). 3. Upper Jurassic (Oxfordian-Tithonian). 4. Aptian flysch. 5. Răciu Breccia (Aptian). 6. Middle Bucegi Conglomerates. 6a. Sandstone facies. 7. Scropoasa-Lăptici Sandstone (Albian). 8. Upper Bucegi Conglomerates and Costila conglomerates (Albian). 9a. Babele Sandstone (Albian). 9b. Sandstone-conglomeratic facies equivalent to Babele Sandstone (Albian). Location of the cross section lines on the index map (left).

Several longitudinal faults (in relation with the general north-south direction of the Albian deposits) from the high plateau and the eastern slope of the Bucegi, were mapped and described by Patruilius (1969) Massif. The Obârşia Mountain, the study area of the present paper, is crossed by the Jepsi Valley Fault, originally identified by Oncescu (1943) and subsequently reconsidered by Patruilius (1969). This fault extends from the eastern slope of the Doamnele Mountain, continues through the south foot of the Obârşia Mountain,

to the upper Jepsi River (Fig. 5). Several short NNE to SW oriented faults are limiting the eastern extension of the Jepsi Valley Fault.

In the southern part of Obârşia Mountain, a longitudinal fault with limited extension which crosses the Jepsi Valley Fault was mapped out by Patruilius (1969). At the north extremity of the Obârşia Mountain, in the upper part of Cerbului River, Patruilius (1969) mapped two small faults (Obârşia Faults). A small graben is outlined between these faults.




Fig. 5. The main fault lines affecting the Albian deposits in the Obârşia Mountain area (traced in red). Picture from the original geologic map by Patruilius (1969). **Legend:** al Cbm - Middle Bucegi Conglomerates. al GB - Babele Sandstone. al Cg - sandy conglomerates. The index map on the left (simplified Bucegi geological map, based on Patruilius, 1969) shows the location of area.

2.3. GEOLOGICAL MAP IN THE OBÂRŞIA MOUNTAIN AREA

In the initial stage (1948-1951) the entire mountainous massif was mapped by Patruilius (unpublished) at the 1:20,000 scale. When aerial photographs became available, in the 1960-1963 interval, Patruilius (1969) revised and completed the entire Bucegi Massif geologic map. In the Obârşia Mountain area, on the geological map of Patruilius (1969), the Babele Sandstone deposits are exposed on the mountain slopes, while the underlying Upper Bucegi Conglomerate deposits crop out along the Şugărilor and Obârşiei Rivers (Fig. 6).

On the two Obârşia Mountain summits, Patruilius (1969) mapped out conglomerates intercalated in deposits of the Babele Sandstone Member. These conglomerates, associated with olistoliths, were named “Obârşia-type conglomerates”, and described in the book chapter (Patruilius, 1969) dealing with the Upper Bucegi Conglomerates.

Two levels of conglomerates are separated by Patruilius (1969) in the northern summit, *i.e.*, the Colţii Obârşiei Peak. The Obârşia-type conglomerates from the southern summit

are mapped as a single body. Another conglomerate body was described south of the southern summit.

3. RESULTS

3.1. LITHOSTRATIGRAPHY

As illustrated by the panoramic view at the Obârşia Mountain slopes (Fig. 7), four conglomerate units associated with olistoliths are intercalated in the upper part of the Babele Sandstone Member. Visible on the satellite image (Fig. 8), the lower conglomerate unit (Unit 1) crops out at “La Cerdac” point (Fig. 9), evidenced as a 25 m high cliff. A 15-20 m thick series of sandstones and microconglomerates occurs between the conglomerate units 1 and 2. At the Colţii Obârşiei peak (the northern summit of the Obârşia Mountain), the units 2, 3 and 4 make up an amalgamated conglomerate succession, with no interbedded Babele Sandstone sediments (Fig. 9). In the area of the southern Obârşia summit, the conglomerate units 2, 3 and 4 occur as separate bodies, with consistent intercalations of Babele Sandstone deposits (Fig. 10). The conglomerate unit 1 was not identified in this area.




Fig. 6. Cartographic image of the Obârșia Mountain area, from the geological map of the Bucegi Mountains by Patrușiu (1969). **Left** - Location of the Obârșia Mountain area, on the simplified geological map of the Bucegi Massif (based on Patrușiu, 1969)




Fig. 7. Conglomerate units (1 to 4) associated with olistoliths, intercalated in the Babele Sandstone Member. View at the eastern side of the Obârșia Mountain, from the tourist path Babele-Omu.

The conglomerates forming the four units series have a poorly sorted, coarse to medium-grained texture (Fig. 11). The sandy matrix is abundant. Large clasts of more than 10 cm diameter and relatively small olistoliths (up to 10-15 m in size) are embedded in the conglomerate mass (Figs. 9 and 10). The conglomerate units 1 and 2 show individual fining upward trend (Fig. 12). At the lower part of the units, there are coarse-grained, massive conglomerates. Well-bedded, very coarse-grained sandstones occur at the top of the fining upward beds. The transition is gradual. The conglomerate units are 15 to 25 m

thick. Their thickness is difficult to determine and most likely is variable. Only the conglomerate unit 1, at "La Cerdac" location (Fig. 9), clearly shows the top and bottom limits, and its thickness was correctly determined (26 m). Usually, the conglomerate unit boundaries are covered (Fig. 10), with only the upper boundary exposed. Poorly cropping out, the Babele Sandstone sediments, which occur between the conglomerate units, consist of medium and coarse-grained sandstones and micro-conglomerates, with cross and flat lamination (Fig. 13).






Fig. 8. Upper Albian conglomerate units associated with olistoliths in the northern part of the Obârșia Mountain. **Left** - blank Bing Maps aerial image. **Right**- conglomerate units outcrop lines traced on the aerial image.






◀ **Fig. 10.** Conglomerate units 2, 3 and 4 at the southern summit of the Obârșia Mountain identified on satellite image (A) and in photograph (B). Babel Sandstone deposits (BS) occur between the three conglomerate units. The upper picture - Bing Map satellite image.



▼ **Fig. 11.** Textural aspects of the conglomerate units from the Obârșia Mountain area (Albian). **A** - Unit 1 conglomerates, at "La Cerdac" point. **B** - Unit 2 conglomerates, close to Colții Obârșiei Peak. **C** - Abundant sandy matrix in the Unit 3 conglomerates, south summit of the Obârșia Mountain. **D** - Large clast in the Unit 4 conglomerates at the south summit of the Obârșia Mountain.






Fig. 12. Fining upward conglomerate beds, evolving from massive rudites to bedded sandstone. **Left** - Unit 1, cropping out in the saddle area between Obârșia and Cerbului rivers. **Right** - Unit 2 conglomerates, western side of the Sugarilor River.




Fig 13. Sandstones and micro-conglomerates with current structures (Babele Sandstone, Albian) occurring below and between the conglomerate units 1 and 2. **A** - Microconglomerates filling a small channel, cut into middle to coarse-grained sandstones. Sediments occurring under the Unit 1 conglomerates. Tourist path to Omu chalet, at "La Cerdac" point. **B** - Medium-scale cross laminated microconglomerates with a lens-shaped sandstone body, covering Unit 1 conglomerates. "La Cerdac" point.

3.2. DETAILED GEOLOGICAL MAPPING

The mapping of the conglomerate units discussed in the present paper was carried out in the area of the two topography summits of the Obârșia Mountain (2488 m and 2385 m high). The outcrop lines of the conglomerate beds are roughly concentric around these heights (Fig. 14).

The outcrop condition of the mapped conglomerate beds is dominantly good or very good, but there are also sites with poor exposure or no outcrops. For a good evaluation of the conglomerate unit cartographic image, the outcrop quality is indicated on the geological map by symbols showing three degrees of confidence (Fig. 14). The high confidence symbol reflects good outcrop coverage of the conglomerate

units and very good aerial image information. Poor outcrop condition or only modest satellite image information are indicated by the moderate confidence sign. The low confidence symbol applies to the absence of data, when the conglomerate beds were tentatively extended on the map between two known points. The field outcrop survey together with satellite images examination (from the Bing Map imagery) (Figs. 8 and 10) indicate that the conglomerates are stratiform and continuous units. The conglomerate bodies do not show significant thickness and grain-size variations across the mapped area. Unlike the conglomerate units 2, 3 and 4, which extend through all the studied area, the conglomerate unit 1 shows considerably smaller extension. This basal unit was traced only in the northeastern part of the Obârșia Moun-




Fig. 14. Detailed geologic map of the stratiform conglomerate units intercalated in the Babel Sandstone deposits (Albian). Obârșia Mountain area, Bucegi Massif. The geological boundary between the Upper Bucegi Conglomerates and the Babel Sandstone after Patrușiu (1969), with amendments. The index map on the left (simplified Bucegi geological map, based on Patrușiu, 1969) shows the location of the map area. The conglomerate units are marked as 1 (lower unit), 2, 3 and 4 (upper unit).

tain, west and south-east of the “La Cerdac” cliff (Fig. 14). The cartographic image (Fig. 14) confirms the compact packing of the conglomerate units 2, 3 and 4, without interbedded sandy deposits, in the Colții Obârșiei Peak area. Thick Babele Sandstone deposits occur between the conglomerate units 2, 3 and 4, a fact well documented by the geologic map (Fig. 14). As it is visible from the geologic cross section (Fig. 15), the conglomerate units are part of a monoclin in the investigation area, with a southward 6-8° apparent slope (general bed dipping is around 20° according to Patrușiu, 1969). The cross section shows the areal continuity of the conglomerate units 2, 3 and 4 and the southward appearance of Babele Sandstone deposits between the conglomerate entities.

3.3. TECTONICS

The conglomerate units system in the Obârșia Mountain area displays a simple structural architecture. No plastic deformation was revealed, but faulting is rather common, and some of the faults affect the conglomerate units setting. On the Obârșia crest, between the two summits, direct observations and aerial or photographic images reveal fault deformation of the conglomerate units 2 and 3 (Fig.16). Three fault lines (F1, F2 and F3), with E-W or ENE-WSW trend, were mapped out. The amount of fault vertical displacement appears to be in the 15-20 m range, according to direct observations (Fig. 16).

Other faults, with similar small vertical displacement, affect older deposits from the study area, showing that faulting was probably postdepositional. Faults were detected at the boundary between the Upper Bucegi Conglomerates and the Babele Sandstone, two members of the Bucegi Formation.

4. DISCUSSION

4.1. THE ALBIAN BUCEGI CONGLOMERATE SUCCESSION IN TERMS OF THE IUGS STRATIGRAPHY NOMENCLATURE

More than one lithostratigraphic names define the different components of the Bucegi conglomeratic succession. For a better, international understanding of the Bucegi conglomeratic

mass, the hierarchy of these lithostratigraphic terms is approached in this paper according to the stratigraphic classification and terminology recommended by the International Subcommission on Stratigraphic Classification of IUGS (Murphy M. A., Salvador, A., Eds., 1999) (Fig. 17).

Initially outlined by Murgeanu and Patrușiu (1963), the complete conglomeratic succession from the Bucegi Massif was presented by Patrușiu (1969). According to this author, the Bucegi stratigraphic pile include *Lower Bucegi Conglomerates* (the basal unit), *Middle Bucegi Conglomerates*, *Scropoasa-Laptici Sandstone*, *Upper Bucegi Conglomerates* and *Babele Sandstone* (the top unit).

The term Conglomerates with Olistoliths is introduced in the present paper. This new name refers to the four conglomerate units with olistoliths intercalated into the Babele Sandstone and the conglomerate mass resulting through the amalgamation of these units, in the northwestern area of the Bucegi Bassif.

Within this Bucegi stratigraphic pile, the Lower Bucegi Conglomerates show distinct characteristics. This unit consists of several relatively large rudaceous bodies occurring within the coarse-grained and thick bedded Upper Aptian sandstones. The stratigraphic position of these bodies is very close to the base of the Middle Bucegi Conglomerates, but, according to Ștefănescu (1980), between the Upper Conglomerates and the top Upper Aptian deposits a nonconformity surface exists.

The succession of the Middle Bucegi Conglomerates, Scropoasa-Laptici Sandstone, Upper Bucegi Conglomerates and Babele Sandstone units make up an assemblage with common lithologic and enetic features. This conglomerate assemblage is unconformity surfaces bounded (Fig. 17), a fact that also reveals its identity. Therefore, the formation rank (Bucegi Conglomerate Formation) can be awarded to the stratigraphic succession of the Middle Bucegi Conglomerates, Scropoasa-Laptici Sandstone, Upper Bucegi Conglomerates and Babele Sandstone units, without including the Lower Bucegi Conglomerates (Fig. 17).




Fig. 15. Geological section showing the conglomerate units intercalated in Babele Sandstone deposits from Obârșia Mountain. The thickness of the conglomerate beds ranges from 15 to 25 m.




Fig. 16. The fault system affecting the conglomerate beds, on the crest of the Obârșia Mountain. **A.** Faults area location on the geological map. **B.** Blank aerial image of the faults area (from Bing Maps). **C.** Conglomerate units (and associated olistoliths) and the faults lines, edited on the aerial image. **D.** The fault lines F1, F2 and F3 marked out on the photograph of the east side of the Obârșia ridge.

Differences exist also between the lithostratigraphic units belonging to the Bucegi Conglomerate Formation. The Middle Bucegi Conglomerates, Upper Bucegi Conglomerates and Babele Sandstone terms describe the main units of this formation. Their extension covers almost the entire Bucegi Mountain area. These lithostratigraphic units are the members of the Bucegi Conglomerate Formation (Fig. 17). The deposits described as Scropoasa–Laptici Sandstone and Conglomerates with Olistoliths represent lateral facies within the Bucegi Conglomerate Formation (Fig. 17).

The IUGS recommended terminology is not in favor of using the terms upper and lower for the formal lithostratigraphic units. This appears to be in contradiction with the terminology of the Bucegi Formation members. The terms Lower, Middle and Upper Bucegi Conglomerates are local lithologic names given by Murgeanu and Patrușiu (1963), which are in use since more than half a century. The attempt

to change these names would have created serious disturbance for the students of the Eastern Carpathians Cretaceous stratigraphy.

4.2. THE MAPPED CONGLOMERATE BEDS – A NEW LITHOSTRATIGRAPHIC UNIT OF THE BUCEGI FORMATION

The investigations carried out during the present study outline a subsidiary conglomeratic stratigraphic unit, of at least 50 m thickness, at the upper part of the Babele Sandstone Member. According to nomenclature of the International Stratigraphic Guide (Murphy M. A. and Salvador, A., Eds., 1999), the four conglomerate bodies of this subsidiary unit will be named “beds”, representing smaller rank lithostratigraphic units included in the Babele Sandstone Member (Fig. 17).

The Bed 1 to Bed 4 series stands for the uppermost strata of the Albian Bucegi Formation. The occurrence of this conglomerate accumulation, associated with olistoliths, grants




Fig. 17. Lithostratigraphic hierarchy and terminology of the conglomeratic Albian deposits in the Bucegi Massif area (Eastern Carpathians), in terms of the IUGS stratigraphy nomenclature. This stratigraphic scheme uses the well-established lithologic units names given by Murgeanu and Patrușiu (1963), and Patrușiu (1969).

a special genetic significance to the northern Babele Sandstone Member, as compared to the sediment accumulation of the same unit in the center and southern part of the massif.

4.4. CORRELATION OF THE CONGLOMERATE UNITS

The geological mapping required rigorous identification of the individual conglomerate units. The units 1 and 4 were usually easy to recognize, as they make the bottom and the top of the conglomerate succession. When they are in direct contact and the cropping out conditions are poor, identification of the units 2 and 3 presents some difficulties. Sedimentary features can be used to overcome the problems.

The conglomerate unit 2 is characteristically fining upward, with continuous transition (Fig. 18A). The conglomerate unit 3 is not graded and shows thin sandstone intercalations (Fig. 18B). The difference is consistent enough to serve for outcrop identification of the two units. The use of these criteria is illustrated by the setting shown in Figure 18C, where the unit 2 directly overlies the unit 3. The limit between the two conglomerate bodies is marked by the sandstone occurrence, representing the top of the fining upward

conglomerate unit 2, while the overlying conglomerate unit 3 shows stratification and no grain-size grading.

4.5. LARGE-SCALE GENETIC FEATURES OF THE CONGLOMERATE UNITS

The geological mapping and its stratigraphic architecture made evident two important characteristics, of high genetic significance for the Obârșia conglomerate units.

The first genetic significant feature is the restricted areal extension of the conglomerate unit 1 and the bypassing of this lower unit by the other three, upper units (Figs. 14 and 15). This is a large scale sedimentary feature documented by field work combined with satellite image examination.

According to our investigation, the conglomerate units 2, 3 and 4, separated by sand deposits (Figs. 10, 14 and 15), amalgamate northwards (at Colții Obârșiei) into a single conglomerate body (Fig. 9 and 15). Farther to the northwest, in the Gaura Valley area, the conglomerate unit 1 also incorporates into this rudaceous body (Conglomerates with Olistoliths). This is the second large-scale characteristic of the conglomerate units assemblage.




Fig. 18. Facies features used as criteria for the conglomerate unit correlation in the Obârșia Mountain area. **A.** Unit 2 is fining upward. Its top sandstone part is stratified and makes extensive and characteristic outcrops. South Obârșia Mountain, eastern side. **B.** Unit 3 shows internal bedding, with thin sandstone intercalations, and is not graded. South Obârșia Mountain, eastern side. **C.** Separation between Unit 2 and the overlying Unit 3 is based on the identification of the sandstone occurring at the top of the fining upward Unit 2. Northern part of Obârșia Mountain, the eastern side.

5. CONCLUSIONS

The Albian conglomerate deposits of the Obârșia Mountain area (northern Bucegi Massif) have been subjected to detailed field-work. The goal of the study was to investigate the stratigraphic and tectonic features of the series of four conglomerates associated with olistoliths, intercalated at the upper part of the Babele Sandstone Member, the youngest lithostratigraphic unit of the Bucegi Conglomerate Formation. The sedimentologic interpretation of the conglomerate units will be dealt with in a forthcoming study.

To the northwest, the four conglomerate units merge into a single conglomerate body. In the present paper, these rudaceous sediments are named “Conglomerates with Olistoliths”. The sedimentogenetic interpretation of these conglomerates will make the subject of a forthcoming study.

The Conglomerates with Olistoliths define a distinct lithostratigraphic assemblage. According to their stratigraphic

position, the Conglomerates with Olistoliths are a lateral facies of the Babele Sandstone unit and overlies the Upper Bucegi Conglomerates (Fig. 17).

ACKNOWLEDGEMENTS

Dr. Mihaela Melinte-Dobrinescu (National Institute of Marine Geology and Geo-ecology) kindly reviewed an earlier version of the manuscript, offering suggestions which improved the stratigraphic interpretation of the field data.

The authors are grateful to Prof. Nicolae Anastasiu (University of Bucharest, Department of Geology and Geophysics), Dr. Cornel Olariu (University of Texas at Austin, Department of Geological Sciences), and Dr. Sorin Radan (Geological Institute of Romania) for their constructive reviews.

This work was supported by a grant of the Romanian National Authority for Scientific Research, CNCS – UEFISCDI, project number PN-II-PCE-2011-3-0162.

REFERENCES

- BRALOWER, T. J., FULLAGAR, P. D., PAULL, C. K., DWYER, G. S., LECKIE, R. M., 1997. Mid-Cretaceous strontium-isotope stratigraphy of deep-sea sections. *Geol. Soc. Am. Bull.* 109, 1421-1442.
- BRICEAG, A., JIPA, D., MELINTE, M. 2009. Early Cretaceous deposits of the Ceahlău Nappe (Romanian Carpathian Bend area). *Geo-Eco-Marina* 15, 177-185.
- DUMITRESCU, I., SÂNDULESCU, M., 1968. Problèmes structuraux fondamentaux des Carpates roumaines et de leur avant-pays. *Annuaire du Comité Géologique* 36, 195-218.
- MELINTE, M.C., JIPA, D., 2007. Stratigraphy of the Lower Cretaceous sediments from the Romanian Carpathian Bend area. *Acta Geologica Sinica, English Edition*, 81, 6, 949-956.
- MURGEANU, G., PATRULIUS, D., 1957. Le Crétacé supérieur de la Leaota et l'âge des Conglomerats de Bucegi. *Rev. Géol. Géogr.* 1, 109-124.
- MURGEANU, G., PATRULIUS, D., 1963. Les conglomérates de Bucegi, formation de molasse mésocrétacée des Carpates Orientales. *Association Association Géologique Carpato-Balkanique. VI-ème Congrès. Résumés des communications*, 113-115.
- MURGEANU, G., PATRULIUS, D., CONTESCU, L., JIPA, D., MIHĂILESCU, N., PANIN, N., 1963. Stratigrafia si sedimentogeneza terenurilor cretacice din partea internă a Curburii Carpaților. *Asociația Geologică Carpato-Balcanică. Congresul V. Comunicări științifice. Secția II stratigrafie III/2*, 31-58.
- MURPHY M. A., SALVADOR, A. (Eds.), 1999. *International Stratigraphic Guide - An abridged edition*. International Subcommission on Stratigraphic Classification (ISSC). *Episodes*, 22/ 4, 255-271.
- MUTTERLOSE, J., 1992. Lower Cretaceous nannofossil biostratigraphy of Northwestern Australia (Leg 123). In: Gradstein, F. M., Ludden, J. N., *et al.* (Eds.), *Proceedings of the Ocean Drilling Program, Scientific Results* 123, 343-368.
- ONCESCU, N., 1943. Région de Piatra Craiului-Bucegi. Étude géologique. *Anuarul Institutului Geologic al României* 22, 1-124.
- PATRULIUS, D., 1969. *Geologia Masivului Bucegi și a Culoarului Dâmbovicioara*. Editura Academiei Române, 321 pp.
- PATRULIUS, D., DIMITRESCU, R., POPESCU, I., 1971. Geological map of Romania, scale 1:50,000, sheet 110d Moeciu. Edited by the Geological Institute of Romania.
- ROTH, P.H., 1983. Jurassic and Lower Cretaceous calcareous nannoplankton in the western North Atlantic (Site 534): biostratigraphy, preservation and some observations on biogeography and palaeoceanography. *Initial Reports of Deep Sea Drilling Projects* 76, 587-621.
- SÂNDULESCU, M., 1984. *Geotectonica României*. Editura Tehnică București, 336 pp.

